

RVB gauge theory and the Topological degeneracy in the Honeycomb Kitaev model

Saptarshi Mandal,¹ R. Shankar,² and G. Baskaran^{2,3}

¹*Theoretical Physics Division, IACS, Jadavpur, Kolkata-700032, India.*

²*The Institute of Mathematical Science, CIT Campus, Chennai-600113, India.*

³*Perimeter Institute for Theoretical Physics, 31 Caroline St N, Waterloo, Ontario, Canada N2L 2Y5*

We relate the Z_2 gauge theory formalism of the Kitaev model to the $SU(2)$ gauge theory of the resonating valence bond (RVB) physics. Further, we reformulate a known^{16–19} Jordan-Wigner transformation of Kitaev model on a torus in a general way that shows that it can be thought of as a Z_2 gauge fixing procedure. The conserved quantities simplify in terms of the gauge invariant Jordan-Wigner fermions, enabling us to construct exact eigen states and calculate physical quantities. We calculate the fermionic spectrum for flux free sector for different gauge field configurations and show that the ground state is four-fold degenerate on a torus in thermodynamic limit. Further on a torus we construct four mutually anti-commuting operators which enable us to prove that all eigenstates of this model are four fold degenerate in thermodynamic limit.

PACS numbers: 75.10.Jm, 03.67.-a, 03.67.Lx, 71.10.Pm

Keywords:

I. INTRODUCTION

Solid state realization of q-bits that do not decohere easily is a challenging task in the field of Quantum Computation. Topological defects in strongly correlated quantum many body systems are protected from decoherence and have been suggested as q-bit candidates^{1–5}. In this context, Kitaev constructed a remarkable two dimensional quantum spin model that exhibits abelian and non-abelian anyons and is exactly solvable for its spectrum^{6,7}. It was later shown that the spin correlation functions are also exactly solvable⁸. This model is also extremely interesting from point of view of frustrated spin models and the physics of the resonating valence bond (RVB) states^{9–13}. It realizes, in an exact fashion, the phenomenon of quantum number fractionization and emergent gauge fields that were conjectured and approximately realized in 2D models for RVB states or quantum spin liquids^{14,15}. It has also been shown that the Jordan-Wigner transformation in this model yields a local fermionic theory^{16–19}. This makes the Kitaev model an important one that warrants further investigation; and no wonder that an extensive body of research^{20–24} has already been carried out exploring its many fascinating aspects. Kitaev showed that the model has a natural formulation in terms of a Majorana fermion interacting with Z_2 gauge fields. The remarkable feature of the Kitaev model is that the gauge fields turn out to be static. This greatly simplifies the dynamics leading to the exact computation of the spectrum and spin-spin correlation functions.

The gauge theory of spin- $\frac{1}{2}$ models has a recent history. It was initially formulated²⁵ in the context of strongly correlated electronic systems such as a spin- $\frac{1}{2}$ Mott insulator, as a way of implementing the single electron occupancy constraint,

$$\sum_{\sigma} \left(c_{i\sigma}^{\dagger} c_{i\sigma} - 1 \right) |\psi\rangle = 0. \quad (1)$$

This equation is the Gauss law constraint for a $U(1)$ lattice gauge theory. It resulted in a strongly interacting $U(1)$ gauge theory formalism of spin- $\frac{1}{2}$ models. Soon it was realized that in spin- $\frac{1}{2}$ systems, the $U(1)$ gauge invariance always implied an $SU(2)$ invariance^{26–28}. The consequent $SU(2)$ gauge theory formalism was found to be useful in the context of relating apparently different mean field solutions of the model. The extended Hilbert space of the spin- $\frac{1}{2}$ system is much smaller than that of normal lattice gauge theory where the gauge degrees of freedom on the links are $SU(2)$ group elements. Consequently, it was shown that the essential physics of the spin- $\frac{1}{2}$ system is captured by a Z_2 gauge theory^{29,30}, where the Z_2 gauge group is the center of the original $SU(2)$ gauge group. The Z_2 gauge theory formalism has been effectively used to bring out the physics of quantum number fractionization in spin- $\frac{1}{2}$ systems¹⁵.

In this paper we follow the route charted out above in the context of the Kitaev model and show that the Z_2 gauge theory can indeed be thought of as the center of the $SU(2)$ gauge theory of RVB theory. This sheds light on Kitaev's assertion⁶ that the model represents the same universality class of topological order as RVB. We examine the degeneracy of states in the system defined with periodic boundary conditions in detail. We show that this degeneracy, which characterises the topological order in the system, arises from the so called large gauge transformations. Namely, gauge field configurations which correspond to the same flux configuration that are not related to each other by local gauge transformations. These topologically distinct gauge field configurations can be labelled by the value of the Wilson loops that wind around the torus in the two different directions. We find these gauge configurations for the ground state and demonstrate the four-fold degeneracy of the ground state by explicitly computing the energies. Further, we generalise this proof to all eigenstates by constructing the operators that generate the large gauge transformations and showing that they do not change the

energy of the system in the thermodynamic limit.

The rest of the paper is organised as follows. Section (II) reviews the Kitaev honeycomb model with its main features and some mathematical notions which are used in later sections. The $SU(2)$ gauge symmetry of the model is reviewed in section (III) where we show that Kitaev's choice of the representation of the spin operators in terms of Majorana fermions amounts to a $SU(2)$ gauge fixing procedure which fixes the gauge upto the center Z_2 gauge transformations. In section (IV), we construct a generalized Jordan-Wigner transformation and show that it is a Z_2 gauge fixing transformation in the Kitaev formalism. We derive the fermionic Hamiltonian and simplify the conserved quantities in terms of gauge invariant Jordan-Wigner operators. In section (V), we explicitly work out the four fold degeneracies of the Kitaev model on a torus. We derive the fermionic spectrum for flux free sector in (VA) and show that it leads to four degenerate ground states in thermodynamic limit. Following this in section (VB), we demonstrate that every eigenstate of the Kitaev model has four fold degeneracy. To this end we derive four mutually anticommuting operators that commute with the hamiltonian in the thermodynamic limit. The minimum dimension of the representation of the 4-dimensional Clifford algebra is 4. Thus we are able to prove that all eigenstates are at least four fold degenerate. We summarize our results in the section (VI).

II. THE KITAEV MODEL

A. The Hamiltonian

The Kitaev model is a spin- $\frac{1}{2}$ system on a honeycomb lattice. The Hamiltonian is

$$H = -J_x \sum_{\langle ij \rangle_x} \sigma_i^x \sigma_j^x - J_y \sum_{\langle ij \rangle_y} \sigma_i^y \sigma_j^y - J_z \sum_{\langle ij \rangle_z} \sigma_i^z \sigma_j^z, \quad (2)$$

where i, j run over the sites of the honeycomb lattice, $\langle ij \rangle_a$, $a = x, y, z$ denotes the nearest neighbor links oriented in the a 'th direction as shown in Fig. 1. We will be working with periodic boundary conditions which are defined as follows. The honeycomb lattice is a triangular lattice with a basis of two sites. The sites of the triangular lattice are given by,

$$\mathbf{R}_{m,n} = m\mathbf{e}_1 + n\mathbf{e}_2, \quad (3)$$

where m, n are integers and $\mathbf{e}_1 \cdot \mathbf{e}_2 = -\frac{1}{2}$, $\mathbf{e}_1 \cdot \mathbf{e}_1 = 1 = \mathbf{e}_2 \cdot \mathbf{e}_2$. The label i of the sites of the honeycomb lattice therefore stands for (m, n, α) where $\alpha = a, b$ is the sub lattice label. The periodic boundary conditions are then defined by,

$$\sigma_{m,n,\alpha}^a = \sigma_{m+M,n+N,\alpha}^a. \quad (4)$$




FIG. 1: Honeycomb lattice and the Kitaev model

B. The conserved quantities

There is a conserved quantity associated with every plaquette of the lattice. If the plaquette is denoted as p and its vertices labelled as shown in Fig. 1, then, following Kitaev's notation, the conserved quantity is,

$$B_p = \sigma_1^x \sigma_2^z \sigma_3^y \sigma_4^x \sigma_5^z \sigma_6^y. \quad (5)$$

We have $B_p^2 = 1$ implying that the B_p s can take values ± 1 . It is clear that any product of the B_p s will also commute with the Hamiltonian. In fact there is a conserved quantity associated with every closed self avoiding loop, C , on the lattice defined the following way. At every site, the path will pass through two of the three bonds that emanate from it. We call these two bonds as the tangential bonds and the third one the normal bond. We associate two tangential vectors at each site, $\hat{\mathbf{t}}_{1i}$ and $\hat{\mathbf{t}}_{2i}$ which are either $\hat{\mathbf{x}}$, $\hat{\mathbf{y}}$ or $\hat{\mathbf{z}}$ according to the direction of the incoming bond and the outgoing bond respectively. We then define a normal vector $\hat{\mathbf{n}}_i$ as

$$\hat{\mathbf{n}}_i \equiv \hat{\mathbf{t}}_{1i} \times \hat{\mathbf{t}}_{2i}. \quad (6)$$

If the sites of C are $i_1, i_2, \dots, i_N$, then the conserved quantity associated with it is,

$$B_C = \prod_{n=1}^N (\hat{\mathbf{n}}_{i_n} \cdot \sigma_{i_n}). \quad (7)$$

It can be checked that,

$$[B_C, H] = 0, \quad B_C^2 = 1. \quad (8)$$

We will call C topologically trivial if it can be written as a product of B_p s. On the torus, we have two loops which wind the torus around in the two directions which cannot be expressed as a product of B_p s. One cannot be obtained from the other by multiplication by B_p s. We will call these two (Wilson)loops W_1 and W_2 .

All the B_p s are not independent due to the identity,

$$\prod_p B_p = 1. \quad (9)$$

Thus there are $N_p - 1$ independent B_p s, where $N_p = MN$ is the number of plaquettes. Together with W_1 and W_2 , we have a total of $N_p + 1$ conserved quantities on the torus. These two loop operators account for the four fold degeneracy on a torus.

III. FROM $SU(2)$ TO Z_2

A. $U(1)$ and $SU(2)$ gauge symmetry

Interacting quantum spin systems often lead to spontaneously broken symmetric states such as a ferro, antiferro or spiral magnetic states. Low energy physics of these ordered states are captured by the well known spin wave approximations, resulting in Goldstone mode type bosonic low energy effective theories³¹. A quantum spin liquid, on the other hand, has no classical long range order. Experiments in LaCuO_4 and other low spin Mott insulators, according to Anderson, indicated possible presence of neutral fermionic excitations in a quantum spin liquid³²⁻³⁴. A theory to describe such a quantum spin liquid or resonating valence bond (RVB) state needed a paradigm shift from spin wave theory. It was also clear that a quantum spin liquid, in view of different possible phase coherence among disordered spin configurations, could offer a variety of quantum spin liquid states to be realized in nature. RVB gauge theory attempted to capture these new possibilities, through an approach involving enlarged Hilbert space and emergent gauge fields in strongly correlated electron systems. Through the work of Wen³⁶ and others it has become clear that there is a plethora of spin liquid phases, characterized by *quantum order and projective symmetry groups*.

The spin- $\frac{1}{2}$ Hilbert space can be realized as the subspace of the Hilbert space of two fermions defined by the constraint,

$$\left(\sum_{\sigma=\uparrow\downarrow} c_{\sigma}^{\dagger} c_{\sigma} - 1 \right) |\psi\rangle = 0, \quad (10)$$

where $c_{\sigma}^{\dagger}$ and c_{σ} are the fermion creation and annihilation operators. As mentioned earlier, Eq. (10) can be looked upon as the Gauss law constraint for a $U(1)$ gauge theory. The LHS of the equation being the generator of the following $U(1)$ gauge transformations on the fermion operators,

$$c_{\sigma} \rightarrow e^{i\Omega} c_{\sigma}, \quad c_{\sigma}^{\dagger} \rightarrow e^{-i\Omega} c_{\sigma}^{\dagger}. \quad (11)$$

The spin operators,

$$S^a = \frac{1}{2} c_{\sigma}^{\dagger} \sigma_{\sigma\sigma'}^a c_{\sigma'}, \quad (12)$$

are then the gauge invariant observables of the theory. σ^a are the Pauli spin matrices.

The single occupancy constraint in the spin- $\frac{1}{2}$ theory implies that a spin- $\uparrow$ hole is the same as a spin- $\downarrow$ particle in the physical space. This can be mathematically expressed as an $SU(2)$ gauge invariance. It is convenient to express this symmetry in terms of a matrix of the fermion operators

$$\Psi \equiv \begin{pmatrix} c_{\uparrow} & -c_{\downarrow}^{\dagger} \\ c_{\downarrow} & c_{\uparrow}^{\dagger} \end{pmatrix}. \quad (13)$$

In terms of this matrix, the spin operators are given by,

$$S^a = \frac{1}{4} \text{tr} \Psi^{\dagger} \sigma^a \Psi. \quad (14)$$

The generators of the $SU(2)$ gauge transformation are given by,

$$\tilde{S}^a = -\frac{1}{4} \text{tr} \Psi \sigma^a \Psi^{\dagger}. \quad (15)$$

The Ψ matrix transforms under the $SU(2)$ spin and $SU(2)$ gauge transformations as,

$$\Psi \rightarrow U_S \Psi U_G^{\dagger}. \quad (16)$$

Where U_S and U_G are $SU(2)$ matrices representing the spin and gauge transformations respectively. It is clear from equations (14), (15) and (16) that the spin operators are gauge invariant and the generators of gauge transformations are spin singlet. The constraint in Eq. (10) is exactly equivalent to the $SU(2)$ Gauss law,

$$\tilde{S}^a |\psi\rangle = 0. \quad (17)$$

Before we close this section, we wish to mention that the above gauge theory formalism offers a possible way to understand quantum spin liquid states as and when they exist. This formalism does not guarantee a simple gauge theory structure at all energy scales in the physics of the problem. It only suggests that in some systems (for some Hamiltonians) at low energy scales there could be emergent gauge fields and interesting consequences of quantum number fractionization, quantum order etc. At high energy scales gauge fields interact and it is no more simple or useful to talk in terms of emergent gauge fields. As we will see soon, the Hamiltonian invented by Kitaev on a honeycomb lattice is very special. It offers static Z_2 gauge fields and makes the Z_2 gauge theory meaningful at all energy scales.

B. Majorana fermions and the Z_2 theory

We can make connection to Kitaev's representation of the spins by writing,

$$c_{\uparrow} = \frac{c_y - ic_x}{2}, \quad c_{\downarrow} = -\frac{c - ic_z}{2}. \quad (18)$$

where c, c_x, c_y and c_z are Majorana fermions. The single occupancy constraint, Eq. (10) reduces to exactly Kitaev's form,

$$cc_x c_y c_z = 1. \quad (19)$$

Kitaev's representation of the spins then get written as,

$$\frac{i}{2} cc_a = S^a - \tilde{S}^a. \quad (20)$$

Note that these three operators are not equal to the gauge invariant spin operators in the extended Hilbert

space but are exactly equivalent to them in the physical Hilbert space. Substituting the expressions in Eq. (20) for the spin operators in the Hamiltonian is then equivalent to adding gauge fixing terms. Since the $SU(2)$ gauge generators are invariant under the Z_2 center of the gauge group, these terms only fix the gauge upto the central Z_2 group represented by $e^{i2\pi\tilde{S}^3}$. The Hamiltonian will therefore continue to have a Z_2 gauge symmetry.

The simple example of a spin- $\frac{1}{2}$ in a magnetic field illustrates these issues. If we take the Hamiltonian to be

$$H = B S^3. \quad (21)$$

then the theory has $SU(2)$ gauge symmetry, the degenerate ground states in the extended space are

$$|GS\rangle = \alpha c_{\downarrow}^{\dagger}|0\rangle + \beta|0\rangle + \gamma c_{\uparrow}^{\dagger}c_{\downarrow}^{\dagger}|0\rangle, \quad (22)$$

for arbitrary α, β and γ , $|\alpha|^2 + |\beta|^2 + |\gamma|^2 = 1$. The last two states in the RHS of Eq. (22) transform as a doublet under the $SU(2)$ gauge symmetry. Gauge averaging therefore projects out the ground state in the physical subspace, namely $c_{\downarrow}^{\dagger}|0\rangle$.

If the Hamiltonian is taken to be,

$$H = iB cc_z = B (S^3 - \tilde{S}^3). \quad (23)$$

The theory has only Z_2 gauge invariance, the degenerate ground state in the extended Hilbert space are,

$$|GS\rangle = \alpha c_{\downarrow}^{\dagger}|0\rangle + \gamma c_{\uparrow}^{\dagger}c_{\downarrow}^{\dagger}|0\rangle. \quad (24)$$

The second state in the above equation transforms non-trivially under the Z_2 gauge transformation. Thus again, under gauge averaging, the ground state in the physical sector is projected out. Thus Kitaev's representation of the spin operators can be interpreted as adding gauge fixing terms to the $SU(2)$ gauge invariant hamiltonian which leave an unbroken (unfixed) Z_2 gauge symmetry.

IV. THE Z_2 GAUGE THEORY OF THE KITAEV MODEL

A. The Hamiltonian

Following Kitaev⁶, we write the hamiltonian in terms of the Majorana fermions,

$$\tilde{H} = \sum_{a=1}^3 \sum_{\langle ij \rangle_a} i c_i u_{\langle ij \rangle_a} c_j, \quad (25)$$

where the link variables are defined as $u_{\langle ij \rangle_a} = i c_{ai} c_{aj}$. It is natural to express them in terms of the bond fermions⁸ defined as,

$$\chi_{\langle ij \rangle_a} \equiv \frac{c_{ai} + i c_{aj}}{2}. \quad (26)$$

The link variables are then given in terms of the occupancy number of the bond fermions,

$$u_{\langle ij \rangle_a} = 2 \chi_{\langle ij \rangle_a}^{\dagger} \chi_{\langle ij \rangle_a} - 1. \quad (27)$$

It is easy to see that,

$$\begin{aligned} u_{\langle ij \rangle_a}^2 &= 1, \\ [u_{\langle ij \rangle_a}, H] &= 0. \end{aligned} \quad (28)$$

Thus the link variables can be interpreted as static Z_2 gauge fields.

It is remarkable that at one shot Kitaev hamiltonian has been solved exactly for the entire many body spectrum ! Infact, two related phenomena occur: i) the Z_2 gauge theory is exact at all energy scales and ii) the enlarged Hilbert space gets decomposed into sectors that are identical gauge copies having the same energy eigen values (Fig. 2). The Hilbert space enlargement does not produce any unphysical state, but only gauge copies. In the standard $U(1)$ RVB gauge theory for a Heisenberg antiferromagnet, for example, it is easy to see how unphysical states are brought in by Hilbert space enlargement. For example, an unphysical state containing M doubly occupied and M empty sites gives spectrum of a Heisenberg antiferromagnet containing 2M missing sites.




FIG. 2: How the enlarged Hilbert space for Kitaev Model become just Gauge Copies.

B. The Jordan-Wigner transformation and Z_2 gauge fixing

A remarkable feature of the Kitaev model is that the Jordan-Wigner transformation yields a local fermionic Hamiltonian^{16–19}. In this section we show that the Jordan-Wigner transformation in the Kitaev model can be interpreted as a Z_2 gauge fixing procedure resulting in gauge invariant (gauge fixed) Majorana and bond fermions. The choice of the Jordan Wigner path amounts to a choice of the Z_2 gauge.

1. The Jordan-Wigner fermionisation

We define the Jordan-Wigner transformation as follows³⁹. Take any Hamilton path on the lattice defined by a sequence of sites i_n , $n = 1, \dots, N_S$, where N_S is the number of sites in the lattice. The path will classify each bond as normal or tangential as defined in section II B. The normal bonds will form a dimer covering of the lattice. For a given site ‘ i ’ we attach three vectors, two tangential vectors denoted by $\hat{\mathbf{t}}_{1i}$, $\hat{\mathbf{t}}_{2i}$ and one normal vectors $\hat{\mathbf{n}}_i$, such that they follow Eq. (6). Then we can define the Jordan-Wigner transformations in the following compact way,

2. Gauge invariant Jordan-Wigner fermions

To define the Jordan-Wigner transformations we associate two Majorana fermions (η_{i_n} and ξ_{i_n}) at a given site ‘ i_n ’. These Jordan-Wigner (Majorana) fermions are defined in terms of the gauge invariant spin-operators in the following way ,

$$\eta_{i_n} = \hat{\mathbf{t}}_{1i_n} \cdot \sigma_{i_n} \mu_{i_n}. \quad (29)$$

$$\xi_{i_n} = \hat{\mathbf{t}}_{2i_n} \cdot \sigma_{i_n} \mu_{i_n}. \quad (30)$$

$$\mu_{i_n} = \prod_{m=1}^{n-1} (\hat{\mathbf{n}}_{i_m} \cdot \sigma_{i_m}). \quad (31)$$

It can be easily checked that the above definitions refer to the usual anti-commutations relations for Majorana fermions,

$$\begin{aligned} \{\eta_i, \xi_j\} &= 2\delta_{ij}, \\ \{\xi_i, \eta_j\} &= 2\delta_{ij}, \\ \{\eta_i, \xi_j\} &= 0. \end{aligned} \quad (32)$$

Since the Jordan-Wigner fermions are constructed entirely from the spin operators, they are manifestly gauge invariant. However, it is also interesting to see this by rewriting equations (30) and (29) in terms of the original Majorana fermions and gauge fields,

$$\eta_{i_n} = i c_{i_n} (u_{i_{n-1}i_{n-2}} \dots u_{i_2i_1}) \hat{\mathbf{t}}_{1i_1} \cdot \mathbf{c}_{i_1}. \quad (33)$$

$$\xi_{i_n} = i c_{i_n} \cdot \hat{\mathbf{n}}_{i_n} (u_{i_{n-1}i_{n-2}} \dots u_{i_2i_1}) \hat{\mathbf{t}}_{1i_1} \cdot \mathbf{c}_{i_1}. \quad (34)$$

The transformation can be inverted to write the spins in terms of the fermions,

$$\hat{\mathbf{n}}_{i_n} \cdot \sigma_{i_n} = i \eta_{i_n} \xi_{i_n}. \quad (35)$$

$$\hat{\mathbf{t}}_{1i_n} \cdot \sigma_{i_n} = \eta_{i_n} \mu_{i_n}. \quad (36)$$

$$\hat{\mathbf{t}}_{2i_n} \cdot \sigma_{i_n} = \xi_{i_n} \mu_{i_n}. \quad (37)$$

These completes the definitions of Jordan-Wigner fermionisations used in this article.

3. The gauge fixed Hamiltonian

The Hamiltonian can be written in terms of the gauge invariant fermions as,

$$\tilde{H} = J_x \sum_{\langle ij \rangle_x} i \eta_i \tilde{u}_{ij} \eta_j + J_y \sum_{\langle ij \rangle_y} i \eta_i \tilde{u}_{ij} \eta_j + J_z \sum_{\langle ij \rangle_z} i \eta_i \tilde{u}_{ij} \eta_j, \quad (38)$$

where the gauge fixed $\mathbb{Z}_2$ fields, $\tilde{u}_{ij}$ are,

$$\begin{aligned} \tilde{u}_{ij} &= i \xi_i \xi_j \quad \text{normal bonds,} \\ &= 1 \quad \text{tangential bonds except } (ij) = (i_1 i_{N_S}), \\ \tilde{u}_{i_1 i_{N_S}} &= \mathcal{S} \\ &= \prod_{n=1}^{N_S} (\hat{\mathbf{n}}_{i_n} \cdot \sigma_{i_n}) \\ &= \left(u_{i_1 i_{N_S}} u_{i_{N_S} i_{N_S-1}} \dots u_{i_3 i_2} u_{i_2 i_1} \right) \prod_{n=1}^{N_S} \eta_{i_n}. \end{aligned} \quad (39)$$

$\mathcal{S}$ is a well known conserved quantity in the one-dimensional applications where it corresponds to the total number of fermions modulo 2 and determines the boundary conditions on the fermions. Thus the Jordan-Wigner transformation is equivalent to a gauge fixing procedure where all the gauge fields on the tangential bonds (except one) are set equal to 1. The choice of the Hamilton path amounts to a gauge choice since it defines which of the bonds are tangential. It also defines the sign in the definition of u_{ij} for the normal bonds. In Eq. (39), the sign corresponds to a Hamilton path which winds regularly in the $\hat{\mathbf{e}}_1$ direction as shown in Fig (3). Hereinafter all explicit computations will be with respect to this path. A general algorithm to go to this gauge, which we will refer to as the Jordan-Wigner gauge, is given in appendix A.




FIG. 3: Jordan-Wigner path on a torus for 4×4 lattice. The numerics at the lattice sites describe how the Jordan-Wigner path traverses the lattice.

4. The fermionic conserved quantities

All the gauge fixed $\mathbb{Z}_2$ fields are conserved quantities. It is convenient to express them in terms of the gauge

fixed bond fermions on the normal bonds,

$$\chi_{ij} \equiv \frac{\xi_i + i\xi_j}{2}, \quad \chi_{ij}^\dagger \equiv \frac{\xi_i - i\xi_j}{2}. \quad (40)$$

The conserved quantities are then the occupation numbers of the bond fermions,

$$\tilde{u}_{ij} = 2 \chi_{ij}^\dagger \chi_{ij} - 1. \quad (41)$$

There are N_p normal bonds and hence the bond fermion occupation numbers form a set of N_p conserved quantities. Thus along with $\mathcal{S}$, we have $N_p + 1$ conserved quantities consistent with the analysis in terms of spin variables in section II B. To see the meaning of $\mathcal{S}$, it is convenient to define complex fermions in the matter sector from the two ξ fermions on every normal bond,

$$\psi_{ij} \equiv \frac{\eta_i + i\eta_j}{2}, \quad \psi_{ij}^\dagger \equiv \frac{\eta_i - i\eta_j}{2}. \quad (42)$$

It can then be shown that,

$$\begin{aligned} \mathcal{S} &= (-1)^{(N_\psi + N_\chi + 1)}, \\ N_\psi &\equiv \sum_{\text{z bonds}} \psi_{ij}^\dagger \psi_{ij}, \\ N_\chi &\equiv \sum_{\text{normal bonds}} \chi_{ij}^\dagger \chi_{ij}. \end{aligned} \quad (43)$$

5. From Kitaev gauge to Jordan-Wigner gauge

We have explained that Jordan-Wigner gauge is a special realisation of Kitaev gauge where all the gauge fields residing on the tangential bonds are fixed to unity. One may wonder whether there exists a gauge transformation on the lattice which renders Eq. (25) to that of Eq. (38). Indeed there exists such a gauge transformations. Referring to the Jordan-Wigner path given in Fig. (3) we do the following gauge transformations at a site ' n '

$$c_n \rightarrow \prod_{i=1, n-1} u_{i, i+1} c_n, \quad 1 < n < N. \quad (44)$$

In the above equation the various indices corresponds to the Jordan-Wigner path in the Fig. (3). $u_{i, i+1}$ s are the Z_2 gauge fields that would normally exists on the link joining site ' i ' and ' $i+1$ ' if we apply the Majorana fermionisation as adopted by Kitaev. After implementing the above gauge tranformations Z_2 gauge fields appear only on the normal bonds. The above gauge transformations yield a new conserve quantity $\mathcal{S}'$ which appear on the bond where the Jordan-Wigner end path meets. The expression for $\mathcal{S}'$ is,

$$\mathcal{S}' = \prod_{i=1, N-1} u_{i, i+1}. \quad (45)$$

The significance of $\mathcal{S}'$ is very similar to $\mathcal{S}$ introduced in Eq. (39).

V. FOUR FOLD DEGENERACY ON A TORUS

Since the gauge fields are static, the problem is reduced to one of non-interacting fermions on a lattice. It is known^{6,37} that the lowest fermionic ground state energy is obtained for the flux free configuration, i.e. $B_p = 1, \forall p$. On the torus, there are four gauge in equivalent configurations for every configurations of B_p as argued in II B. These correspond to the four values of the gauge invariant conserved quantities W_1 and W_2 . First We examine these four different gauge field configurations for flux free sector and compute the corresponding fermionic ground state. Following this we show that this leads to four fold degeneracy of ground states in thermodynamic limit. Next we demonstrate explicitly how to obtain the four fold degeneracy for every configurations of fluxes. To this end we derive the required operators which enables us to obtain any one of the inequivalent gauge field configurations from the other for arbitrary flux configurations.

A. Degenerate ground states on a Torus

To start with we briefly recapitulate the notions of Jordan-Wigner transformation and refer to Fig. (3). The normal bonds are the ones that form the basis of the triangular lattice except for the $(0, n, \alpha)$ line. On this line the normal bonds are the ones between $(0, n, a)$ and $(0, n+1, b)$. We choose the first site of the path to be $i_1 = (0, 0, b)$. The four flux free configurations are then explained as follows. To this end we write the exact Hamiltonian and B_p for this particular realisation of Jordan-Wigner transformation. We divide the Hamiltonian in three parts, H_{int} , H_{bound} and H_{end} . H_{int} includes all the internal bonds and H_{bound} includes all the boundary bonds except one where the Jordan-Wigner end points meets. H_{end} includes the interaction for the bond where the Jordan-Wigner end points meet each other. Similarly all the B_p 's are categorized in the above three different way. Below we write the various parts of the Hamiltonian and B_p 's.

$$\begin{aligned} H_{int} &= \sum_{m,n} iJ_x \eta_{m,n}^a \eta_{m+1,n+1}^b + \sum_{m,n} iJ_y \eta_{m,n}^a \eta_{m,n+1}^b \\ &\quad + \sum_{m,n} iJ_z \tilde{u}_{m,n} \eta_{m,n}^a \eta_{m,n}^b, \end{aligned} \quad (46)$$

where $\tilde{u}_{m,n}$ is defined on each internal z-bonds.

$$\begin{aligned} H_{bound} &= \sum_{m,n} iJ_y \tilde{u}_{m,n} \eta_{m,n+1}^a \eta_{m,n+1}^b \\ &\quad + \sum_{m,n} J_z i \eta_{m,n}^a \eta_{m,n}^b. \end{aligned} \quad (47)$$

Where $\tilde{u}_{m,n}$ is defined on each boundary y-bond. The Hamiltonian for the end bond is given by,

$$H_{end} = -S i \eta_{M,N}^a \eta_{M,N}^b. \quad (48)$$

Now with the definition of ψ fermion and χ fermion we get the equations (46), (47) to be rewritten as,

$$\begin{aligned} H_{int} = & \sum_{m,n} J_x (\psi_{m,n}^\dagger + \psi_{m,n}) (\psi_{m+1,n+1}^\dagger - \psi_{m+1,n+1}) \\ & + J_y (\psi_{m,n}^\dagger + \psi_{m,n}) (\psi_{m,n+1}^\dagger - \psi_{m,n+1}) \\ & + J_z \tilde{u}_{m,n} (2\psi_{m,n}^\dagger \psi_{m,n} - 1). \end{aligned} \quad (49)$$

The Hamiltonian for the slanting bonds,

$$\begin{aligned} H_{bound} = & \sum_{m,n} J_y \tilde{u}_{m,n} (\psi_{m,n}^\dagger + \psi_{m,n}) (\psi_{m,n+1}^\dagger - \psi_{m,n+1}) \\ & + J_z (2\psi_{m,n}^\dagger \psi_{m,n} - 1). \end{aligned} \quad (50)$$

Lastly the Hamiltonian term for the end bond where the end points of the Jordan-Wigner path meet each other is given by,

$$H_{end} = -\mathcal{S} J_z (2\psi_{M,N}^\dagger \psi_{M,N} - 1). \quad (51)$$

Following equations (39), (30) and (42), we rewrite the complete expressions for various conserved quantities appeared in the final form of the Hamiltonian. The conserve quantity $\tilde{u}_{m,n}$ defined on each internal z-link is given by,

$$\tilde{u}_{m,n} = (2\chi_{m,n}^\dagger \chi_{m,n} - 1). \quad (52)$$

Similarly the conserved quantity defined on each boundary y-bonds (which are labeled by the z-bonds it is connected with (i.e $\tilde{u}_{m,n+1}$)) is given by,

$$\tilde{u}_{m,n+1} = (2\chi_{m,n}^\dagger \chi_{m,n+1} - 1). \quad (53)$$

And the $\mathcal{S}$, for the Jordan-Wigner gauge, is given by,

$$\mathcal{S} = -(-1)^{MN+N_\psi+N_\chi}. \quad (54)$$

If $\mathcal{P}^G$ and $\mathcal{P}^M$ denote the parity operators for the gauge fermions and the matter fermions respectively, then we can write $\mathcal{S} = -(-1)^{MN}\mathcal{P}^M\mathcal{P}^G$. From the Fig. 3, we see that a single hexagon always contains two normal bonds where each normal bond is associated with a conserved(static) Z_2 gauge field. B_p for any plaquette is the product of these two conserved Z_2 gauge fields. Thus, $B_p = \tilde{u}_{ij}\tilde{u}_{kl}$, where 'ij' and 'kl' are the normal bonds for the plaquette 'p'. For the end plaquette where Jordan-Wigner path terminates B_p is given by $B_p = -\mathcal{S}\tilde{u}_{ij}\tilde{u}_{kl}$. Now we are in a position to show the ground state degeneracy in thermodynamic limit. To this end we explicitly write the four in-equivalent gauge field configurations corresponding the flux free configurations and write down the corresponding fermionic Hamiltonian. Finally we find the spectra for each of these four fermionic Hamiltonian.

1. Choice 1

Here the flux free configuration is obtained by making all the $\tilde{u}$'s to be 1 and $\mathcal{S} = -1$. The loop conserve quantities are having the following eigenvalue, $W_1 = 1$ and $W_2 = 1$. These particular choice makes the resulting fermionic Hamiltonian transitionally invariant and usual Periodic boundary condition in both the direction can be used to diagonalize the Hamiltonian. We will explicitly write the Hamiltonian in terms of complex fermion ψ . However one can equivalently work in terms of η , Majorana fermion representation. To keep this in mind we will continue to mention appropriate gauge transformations for η fermions as well as ψ fermions. The complete translational invariant Hamiltonian is given by,

$$\begin{aligned} H = & \sum_{m,n} J_x (\psi_{m,n}^\dagger + \psi_{m,n}) (\psi_{m+1,n+1}^\dagger - \psi_{m+1,n+1}) \\ & + J_y ((\psi_{m,n}^\dagger + \psi_{m,n})) (\psi_{m,n+1}^\dagger - \psi_{m,n+1}) \\ & + J_z (2\psi_{m,n}^\dagger \psi_{m,n} - 1). \end{aligned} \quad (55)$$

This is a manifestly p-wave superconducting Hamiltonian and can be easily diagonalized by going to the momentum space. The constraint on the number of ψ fermion becomes,

$$\prod (2\psi_{m,n}^\dagger \psi_{m,n} - 1) = 1, \quad (56)$$

which implies that we are to fill up only the even number of ψ fermions. We define the Fourier transform of the $\psi_{m,n}$ as given below,

$$\psi_{m,n} = \frac{1}{\sqrt{MN}} \sum_{p,q} e^{i(k_1 m + k_2 n)}, \quad (57)$$

where $k_1 = 2\pi \frac{p}{M}$ and $k_2 = 2\pi \frac{q}{N}$. This is obtained by noticing the fact that we can write $\vec{k} = \frac{p}{M}\vec{G}_1 + \frac{q}{N}\vec{G}_2$ where $\vec{G}_{1/2}$ are the reciprocal lattice vectors which are given by,

$$\vec{G}_1 = \frac{4\pi}{\sqrt{3}} \left(\frac{\sqrt{3}}{2} \mathbf{e}_x + \frac{1}{2} \mathbf{e}_y \right); \quad \vec{G}_2 = \frac{4\pi}{\sqrt{3}} \mathbf{e}_y. \quad (58)$$

Substituting this we get the resulting Hamiltonian in momentum space as,

$$\begin{aligned} H = & \sum_k (\epsilon_k \psi_k^\dagger \psi_k - \epsilon_k \psi_{-k}^\dagger \psi_{-k} + i\delta_k \psi_k^\dagger \psi_{-k}^\dagger - i\delta_k \psi_{-k} \psi_k) \\ & + \epsilon_{0,0} \psi_{0,0}^\dagger \psi_{0,0} + \epsilon_{\pi,0} \psi_{\pi,0}^\dagger \psi_{\pi,0} + \epsilon_{0,\pi} \psi_{0,\pi}^\dagger \psi_{0,\pi} \\ & + \sum_k \epsilon_k - MN J_z. \end{aligned} \quad (59)$$

Where $\epsilon_k = 2(J_x \cos k_x + J_y \cos k_y + J_z)$ and $\delta_k = 2(J_x \sin k_x + J_y \sin k_y)$. $k_x = \mathbf{k} \cdot \mathbf{n}_x$, $k_y = \mathbf{k} \cdot \mathbf{n}_y$ and $\mathbf{n}_{x,y} = \frac{1}{2} \mathbf{e}_x \pm \frac{\sqrt{3}}{2} \mathbf{e}_y$ are unit vectors along x and y type bonds.

In Eq. (59), the sum over 'k' runs over first half of the Brillouin zone and does not include the 'k'-points

$(\pi, 0), (0, \pi), (0, 0)$. The first line of the Hamiltonian is diagonalize by the following transformations,

$$\begin{pmatrix} \alpha_k \\ \beta_k \end{pmatrix} = \begin{pmatrix} \cos\theta_k & -i \sin\theta_k \\ -i \sin\theta_k & \cos\theta_k \end{pmatrix} \begin{pmatrix} \psi_k \\ \psi_{-k}^\dagger \end{pmatrix}. \quad (60)$$

Where $\cos 2\theta_k = \epsilon_k/E_k$, with $E_k = \sqrt{\epsilon_k^2 + \delta_k^2}$. Then re-witting the Hamiltonian we get,

$$\begin{aligned} H = & \sum_k E_k (\alpha_k^\dagger \alpha_k - \beta_k^\dagger \beta_k) \\ & + \epsilon_{0,0} \psi_{0,0}^\dagger \psi_{0,0} + \epsilon_{\pi,0} \psi_{\pi,0}^\dagger \psi_{\pi,0} + \epsilon_{0,\pi} \psi_{0,\pi}^\dagger \psi_{0,\pi} \\ & - \frac{1}{2} (\epsilon_{0,0} + \epsilon_{0,\pi} + \epsilon_{\pi,0}) + \left(\sum_{k'} \frac{1}{2} \epsilon_{k'} - N_z J_z \right) \end{aligned} \quad (61)$$

Here sum over k' runs over full Brillouin zone. The last term in the parenthesis is always zero for Torus.

2. Choice 2

Here the flux free configuration is obtained by making all $\tilde{u}$'s -1 and $\mathcal{S} = -1$. The corresponding values of loop conserve quantity is given by, $W_1 = -1$ and $W_2 = -1$. To implement the Fourier transformation we do the following steps. We make the following gauge transformation. $\eta_{M,n}^b = -\eta_{M,n}^b$ for all ' n '. In terms of ψ fermion the necessary gauge transformation is $\psi_{M,n} = -\psi_{M,n}^\dagger$ for all ' n '. Then the Hamiltonian requires $\eta_{M+1,n} = -\eta_{1,n}$ and $\eta_{m,N+1} = \eta_{m,1}$ (or alternatively $\psi_{M+1,n} = -\psi_{1,n}$ and $\psi_{m,N+1} = \psi_{m,1}$). This is equivalent to anti-periodic boundary Condition in $\mathbf{e}_1$ direction and periodic boundary condition in $\mathbf{e}_2$ direction. The necessary Fourier transform is defined with,

$$\psi_{m,n} = \frac{1}{\sqrt{MN}} \sum_{p,q} e^{i(k_1 m + k_2 n)}, \quad (62)$$

with $k_1 = \frac{2\pi}{M}(m + \frac{1}{2})$; $k_2 = 2\pi \frac{n}{N}$. Substituting this in the Hamiltonian and diagonalizing straightforwardly we get,

$$H = \sum_k E_k^2 (\alpha_k^\dagger \alpha_k - \beta_k^\dagger \beta_k) + \left(\sum_{k'} \frac{1}{2} \epsilon_{k'} - N_z J_z \right) \quad (63)$$

Here also ' k ' runs over first half of the Brillouin zone and the ' k ' runs over the full Brillouin zone. Note the absence of $(0, \pi), (\pi, 0), (0, 0)$ mode. They do not appear here for this anti-periodic boundary condition. This will be true for the choices 3 and 4 also. Various parameters appearing in Eq. (63) are given below,

$$\begin{aligned} E_k^2 &= \sqrt{(\epsilon_k^2 + \delta_k^2)}, \\ \epsilon_k &= 2(J_x \cos k_x + J_y \cos k_y - J_z), \\ \delta_k &= 2(J_x \sin k_x + J_y \sin k_y). \end{aligned} \quad (64)$$

3. Choice 3

For this case the flux free configuration is obtained by making $\tilde{u}_{M,n} = -1$ where ' n ' runs from 1 to $N-1$. All other $\tilde{u}$'s are 1 and $\mathcal{S} = 1$. The loop conserve quantity W_1 takes value 1 and $W_2 = -1$. Similar to the previous case we need to do following gauge transformations in order to apply Fourier transform. We make $\eta_{m,N}^b = -\eta_{m,N}^b$ for all ' m '. In terms of ψ fermion the necessary gauge transformation is $\psi_{m,N} = -\psi_{m,N}^\dagger$ for all ' m '. The resulting Hamiltonian requires $\eta_{m,N+1} = -\eta_{m,1}$ and $\eta_{M+1,n} = \eta_{1,n}$ ($\psi_{m,N+1} = -\psi_{m,1}$ and $\psi_{M+1,n} = \psi_{1,n}$). This indicates anti-periodic boundary condition in $\mathbf{e}_2$ direction and periodic boundary condition in $\mathbf{e}_1$ direction. The resulting Fourier transform is defined with,

$$k_1 = 2\pi \frac{m}{M}; \quad k_2 = \frac{2\pi}{N}(n + \frac{1}{2}). \quad (65)$$

The resulting Hamiltonian in ' k ' space is similar to Eq. (63) with $\epsilon_k = 2(J_x \cos k_x + J_y \cos k_y + J_z)$.

4. Choice 4

For this choices we need, $\tilde{u}_{M,n} = 1$ where ' n ' runs from 1 to $N-1$. All other $\tilde{u}$'s are -1 and $\mathcal{S} = 1$. $W_1 = -1$ and $W_2 = 1$. In this case one requires combined gauge transformation mentioned for the choices 2 and 3. This makes the Hamiltonian anti periodic in both the directions. The required Fourier transform is defined with,

$$k_1 = \frac{2\pi}{M}(m + \frac{1}{2}); \quad k_2 = \frac{2\pi}{N}(n + \frac{1}{2}). \quad (66)$$

Proceeding as before we get exactly Eq. (63) with identical expression for ϵ_k .

5. Ground State Energy in thermodynamic limit

To get the ground state energy for the gauge choices 2,3 and 4, one fills up the negative energy states consistent with the boundary condition (i.e, to satisfy the constraint $\mathcal{S}$ which restricts the total number of particles (odd or even number) to be taken). In the limit of $M, N \rightarrow \infty$ all the above four choices, the ground state energy is obtained as,

$$E_G = \frac{\sqrt{3}}{16\pi^2} \int_{BZ} E(k_x, k_y) dk_1 dk_2, \quad (67)$$

with $E(k_x, k_y)$ is defined before. The appearance of a '-' sign in the expression of ϵ_k for choices 2 and 4 can be accounted for by shifting the k_1 integral to $\pi - k_1$. Thus it is clear that in thermodynamic limit ground state has four fold degeneracy.

B. Four fold Topological degeneracy for any eigenstate

In the preceding section, we have shown that four degenerate ground states are characterized by four different topologically distinct gauge field configurations. From the section IV, we infer that every flux configuration is characterized by such four topologically distinct gauge field configurations. This leads to a four-fold degeneracy for every eigenstate, including the ground state, in the thermodynamic limit.

We now demonstrate this explicitly. Our proof is similar to that of Wen and Niu³⁸ which shows the topological degeneracy of fractional quantum Hall states on a torus. We construct two operators which we call V_1 and V_2 , that act on states with a given gauge field configuration and produce states with a different gauge field configuration without changing the values of the flux operators, B_p . They however change the values of the Wilson loop operators, $W_{1,2}$. These two operators are therefore the generators of the so called “large gauge transformations”. There are four topologically different sectors of gauge field configurations corresponding to $W_i = \pm 1$. We further show that only effect of the large gauge transformations on the matter sector is to change the boundary conditions of the Majorana fermions from periodic to anti-periodic or vice versa. Thus the energy eigenvalues only change by $\sim 1/L$, where L is the length of the torus. The eigenstates in the four sectors (related by the action of V_i) are therefore degenerate in the thermodynamic limit.

The four operators V_i , W_i characterise the topological degeneracy. We show that V_i and W_i satisfy the following algebra,

$$\{V_1, V_2\} = 0, \quad \{V_i, W_i\} = 0, \quad [V_i, W_j]_{i \neq j} = 0. \quad (68)$$

We can then construct four operators, T_a , $a = 1, \dots, 4$,

$$T_1 = V_1 W_1, \quad T_2 = V_2 W_2, \quad T_3 = V_1, \quad T_4 = V_2. \quad (69)$$

These satisfy the Clifford algebra,

$$\{T_a, T_b\} = 2\delta_{ab} \quad (70)$$

Thus we show that the four-fold topological degeneracy on the torus is characterised by the 4-dimensional Clifford algebra.

We will first write down and discuss the expressions for V_1 and V_2 for the 32 site system illustrated in Fig. 3. The construction easily generalises for any even-even lattice.

$$V_1 = \sigma_1^z \sigma_3^z \sigma_5^z \sigma_7^z \quad (71)$$

$$V_2 = \sigma_1^y \sigma_{16}^y \sigma_{17}^y \sigma_{32}^y \prod_{i=2}^7 \sigma_i^z \prod_{i=18}^{23} \sigma_i^z \quad (72)$$

The two Wilson loops for this lattice are,

$$W_1 = \sigma_1^y \sigma_8^y \sigma_9^y \sigma_{16}^y \sigma_{17}^y \sigma_{24}^y \sigma_{25}^y \sigma_{32}^y \quad (73)$$

$$W_2 = \prod_{i=1}^8 \sigma_i^z \quad (74)$$

It can be verified that the above constructions satisfy the algebra in equation (68) and hence the topological operators defined in equation (69) satisfy the Clifford algebra (70). It can also be verified that V_i commute with all the B_p 's.

Now consider simultaneous eigenstates of the Wilson loop operators,

$$W_i |w_1, w_2\rangle = w_i |w_1, w_2\rangle, \quad w_i = \pm 1 \quad (75)$$

The algebra in equation (68) implies,

$$V_1 |w_1, w_2\rangle = |-w_1, w_2\rangle, \quad V_2 |w_1, w_2\rangle = |w_1, -w_2\rangle \quad (76)$$

Thus we have shown that V_i are the generators of large gauge transformations.

We next consider their action on the hamiltonian.

$$V_1 H V_1^{-1} = H^1, \quad V_2 H V_2^{-1} = H^2 \quad (77)$$

where H^1 is the same as H , except that the bonds on one non-trivial loop in the $\mathbf{e}_1$ direction have changed sign, namely the loop $(1, 2, 3, 4, 5, 6, 7, 8, 1)$. In H^2 , a line of parallel bonds in the $\mathbf{e}_1 + \mathbf{e}_2$ direction have changed sign. Namely, the bonds $(7 - 8)$, $(31 - 32)$, $(23 - 24)$ and $(16 - 17)$. We will now write down the operators for a general even-even lattice and then show that the transformed hamiltonians are degenerate in the thermodynamic limit. We will show that in the fermionised theory, these changes of sign can be absorbed into the single particle eigenfunctions of the Majorana fermions and change the single particle energy eigenvalues by $\sim 1/L$. Thus the energies do not change in the thermodynamic limit, making every many body eigenstate four-fold degenerate.

The general expressions for $V_{1(2)}$ are given by

$$V_1 = \prod_{m=1}^M \sigma_{m,0}^{b,z}, \quad V_2 = \prod_{n=1}^{N/2} \sigma_{0,2n}^{a,y} \sigma_{0,2n}^{b,y} \prod_{n=1, N/2}^{m=1, M-1} \sigma_{m,2n-1}^{a,z} \sigma_{m,2n-2}^{b,z}. \quad (78)$$

It can be verified that these constructions satisfy the algebra in equation (68) and also commute with all the B_p 's. In H^1 , the bonds $(m, 0, b) - (m+1, 1, a)$ and $(m, 1, a) - (m, 0, b)$ change sign and in H^2 , the bonds $(M-1, n, b) - (0, n+1, a)$ change sign.

In the fermionised theory, the single particle eigenfunctions satisfy the equation,

$$\sum_j i A_{ij} \phi_j^n = \epsilon^n \phi_i^n \quad (79)$$

where A_{ij} is an antisymmetric matrix coupling the nearest neighbours of the honeycomb lattice. The eigenvalues come in pairs and we denote $(\phi^n)^* = \phi^{-n}$, $\epsilon^{-n} = -\epsilon^n$. n will then go from $1, \dots, NM$. The Hamiltonian is diagonal in terms of the complex fermions defined by,

$$\alpha_n = \sum_i \phi_{ia}^n \eta_i^a + \phi_{ib}^n \eta_i^b \quad (80)$$

$$H = \sum_n \epsilon^n (2\alpha_n^\dagger \alpha_n - 1) \quad (81)$$

We now make the transformation,

$$\phi_{nm,a}^{n'} = e^{in\pi/N} \phi_{nm,a}^n \quad (82)$$

Equation (79) then gets written as,

$$\sum_j A_{ij} \phi_j^n = \sum_j (iA_{ij}^1 + i\delta A_{ij}) \phi_i^{n'} = \epsilon^n \phi_i^{n'} \quad (83)$$

where A^1 is the antisymmetric matrix corresponding to H^1 and $\delta A_{ij} \propto 1/N$ when N is very large. Thus the single particle energy eigenvalues of A and A^1 are identical in the thermodynamic limit when $N \rightarrow \infty$. The spectrum of H and H^1 are also therefore identical. The mapping of the eigenvalues of H and H^2 can be similarly shown using the transformation,

$$\phi_{nm,a}^{n'} = e^{im\pi/M} \phi_{nm,a}^n \quad (84)$$

We can also write equation (78) in terms of the gauge invariant fermions. V_1 or V_2 can be written as $V_i = V_i^{\mathcal{M}} V_i^{\mathcal{G}}$ ($i = 1, 2$). As the notation suggests, $V_i^{\mathcal{M}}$'s and $V_i^{\mathcal{G}}$'s are composed of operators which belong to the matter sector and the gauge sector respectively. They are given by,

$$\begin{aligned} V_1^{\mathcal{G}} &= \prod \xi_{m,0}^a, \quad m = 1, M-1, \\ V_1^{\mathcal{M}} &= \prod \eta_{m,0}^a, \quad m = 1, M, \\ V_2^{\mathcal{G}} &= \prod_{n=1}^{N/2} \xi_{0,2n}^a \xi_{0,2n}^b \prod_{k=2,N}^{j=1,N/2} \xi_{m,2n-1}^a \xi_{m,2n-2}^b, \\ V_2^{\mathcal{M}} &= \prod_{n=1}^{N/2} \eta_{0,2n}^a \eta_{0,2n}^b \prod_{n=2,N/2}^{m=1,M-1} \eta_{m,2n-1}^a \eta_{m,2n-2}^b \end{aligned} \quad (85)$$

From the above expressions we can easily find the commutation relations of parity operator for gauge fermions and the matter fermions. We find that,

$$\{\mathcal{P}^{\mathcal{G}}, V_1\} = 0, [\mathcal{P}^{\mathcal{G}}, V_2] = 0, [\mathcal{P}^{\mathcal{M}}, V_1] = 0, [\mathcal{P}^{\mathcal{M}}, V_2] = 0. \quad (86)$$

The fact that the parity of the matter fermions are conserved is consistent with the fourfold degeneracy discussed here and in (V A).

VI. DISCUSSION

In this paper we have discussed many important aspects of Kitaev model. We have shown how the $SU(2)$ gauge contained in the Kitaev model and explained explicitly how this $SU(2)$ gauge symmetry is reduced to Z_2 gauge symmetry. We have solved the Kitaev model using Jordan-Wigner method in a general way. Though Jordan-Wigner transformations has been used earlier to solve for Kitaev model, our formalism reveals many new features, for example, our definition of J-W transformation is applied to a torus. We focused on the gauge field contents of the Kitaev model and the issue of topological degeneracy. We showed that ground state is four fold degenerate on the torus *in both phases*. While it indicates non-trivial topological order, the ground state degeneracy does not distinguish between the gapless and gapped phases. Finally we have shown the equivalence between the fermionised Hamiltonian obtained in Kitaev gauge and J-W gauge. Lastly we have constructed four mutually anti-commuting operators on a torus to illustrate explicitly the four fold degeneracy for every eigenstate. Our analysis reveals that Jordan-Wigner analysis can be used even in quantum spin liquid problems, as a general method, to bring out non trivial gauge field content, thereby providing an alternative method in resonating valence bond theories.

VII. ACKNOWLEDGEMENTS

As we were completing our manuscript, we learned of a very recent paper by F. J. Burnell and Chetan Nayak (Ref.⁴⁰), which has some overlap with the present paper. GB thanks F. J. Burnell for reference[40]. This research was partly supported by Perimeter Institute for Theoretical Physics.

Appendix A: The gauge fixing algorithm

In this appendix we detail the gauge fixing algorithm to go from any gauge field configuration to one in the Jordan-Wigner gauge. First we will write Hamiltonian as obtained from the Eq. (38). We notice that for $m \neq 0$,

$$\begin{aligned} \hat{\mathbf{t}}_{1m,n}^a &= \hat{y}; \quad \hat{\mathbf{t}}_{2m,n}^a = \hat{x}; \quad \hat{\mathbf{n}}_{1m,n}^a = -\hat{z}, \\ \hat{\mathbf{t}}_{1m,n}^b &= \hat{x}; \quad \hat{\mathbf{t}}_{2m,n}^b = \hat{y}; \quad \hat{\mathbf{n}}_{1m,n}^b = \hat{z}. \end{aligned} \quad (A1)$$

And for $m = 0$,

$$\begin{aligned} \hat{\mathbf{t}}_{1m,n}^a &= \hat{z}; \quad \hat{\mathbf{t}}_{2m,n}^a = \hat{x}; \quad \hat{\mathbf{n}}_{1m,n}^a = -\hat{y}, \\ \hat{\mathbf{t}}_{1m,n}^b &= \hat{x}; \quad \hat{\mathbf{t}}_{2m,n}^b = \hat{z}; \quad \hat{\mathbf{n}}_{1m,n}^b = -\hat{y}. \end{aligned} \quad (A2)$$

Then starting with the Ferromagnetic Hamiltonian we get, for $m = 0, n = 0$,

$$-\sigma_{0,0}^{az} \sigma_{0,0}^{bz} = -i\eta_{0,0}^a \eta_{0,0}^b \mathcal{S}. \quad (A3)$$

For $m \neq 0$

$$-\sigma_{m,n}^{az}\sigma_{m,n}^{bz} = -i\eta_{m,n}^a\eta_{m,n}^b(i\xi_{m,n}^a\xi_{m,n}^b). \quad (\text{A4})$$

now for $m = 0, n \neq 0$

$$-\sigma_{0,n}^{az}\sigma_{0,n}^{bz} = i\eta_{0,n}^a\eta_{0,n}^b. \quad (\text{A5})$$

The above three equations gives complete description of all the z-z interaction. Now for the y-bond we get for $m = 0$

$$-\sigma_{0,n}^{ay}\sigma_{0,n+1}^{by} = -i\eta_{0,n}^a\eta_{0,n+1}^b(i\xi_{0,n}^a\xi_{0,n+1}^b). \quad (\text{A6})$$

For $m \neq 0$ we get

$$-\sigma_{m,n}^{ay}\sigma_{m,n+1}^{by} = i\eta_{m,n}^a\eta_{m,n+1}^b. \quad (\text{A7})$$

At last we write Hamiltonian for x-interaction. For $m = 0$

$$-\sigma_{0,n}^{ax}\sigma_{1,n+1}^{bx} = -i\eta_{0,n}^a\eta_{1,n+1}^b. \quad (\text{A8})$$

And for $m \neq 0$ we get,

$$-\sigma_{m,n}^{ax}\sigma_{m+1,n+1}^{bx} = -i\eta_{m,n}^a\eta_{m+1,n+1}^b. \quad (\text{A9})$$

Now we make the following gauge transformation which we call Jordan-Wigner gauge.

$$\eta_{m,n}^r \rightarrow (-1)^m \eta_{m,n}^r. \quad (\text{A10})$$

Then we choose for each normal bond $\tilde{u}_{i,j} = -i\xi_i^a\xi_j^b$. This gives the Jordan-Wigner Hamiltonian given by

equations 46, 47 and 48. Then we define complex fermion $\chi_{m,n}$ on each normal internal z-bond in the following way,

$$\xi_{m,n}^a = (\chi_{m,n} + \chi_{m,n}^\dagger); \quad \xi_{m,n}^b = \frac{-1}{i}(\chi_{m,n} - \chi_{m,n}^\dagger). \quad (\text{A11})$$

Similarly on each normal slanted y-bond which is joined with a z-link (m, n) and $(m, n+1)$ we define,

$$\xi_{m,n+1}^a = (\chi_{m,n+1} + \chi_{m,n+1}^\dagger); \quad \xi_{m,n+1}^b = \frac{-1}{i}(\chi_{m,n+1} - \chi_{m,n+1}^\dagger). \quad (\text{A12})$$

With this we have alway $\tilde{u}_{i,j} = (2\chi_{i,j}^\dagger\chi_{i,j} - 1)$. At last on each z-link we define ψ in the following way,

$$\eta_{m,n}^a = (\psi_{m,n} + \psi_{m,n}^\dagger); \quad \eta_{m,n}^b = \frac{1}{i}(\psi_{m,n} - \psi_{m,n}^\dagger). \quad (\text{A13})$$

Now the quantity $\mathcal{S}$ for lattice of dimension (M, N) , is given by, $\mathcal{S} = -(-1)^{MN+N_\psi+N_\chi}$. Here N_ψ and N_χ are the number of ψ and χ fermions respectively. Now noticing the number of gauge transformations needed for various flux free configuration we find that we need to fill even number of ψ fermions for each different gauge choices. All the results derived here are based on a representative lattice of dimensions M and N where M and N are both even. It is straight forward to carry the analogous calculation for lattice where M and N can be anything, odd or even. However the results obtained here should not change in thermodynamic limit.

-
- ¹ M. A. Nielsen and I. L. Chang, *Quantum Computation and Quantum Information* (Cambridge University Press, Cambridge, England 2000).
 - ² C. H. Bennett and D. P. DiVincenzo, *Nature* **404** 247(2000).
 - ³ A. Yu. Kitaev, A. H. Shen and M. N. Vyalyi, *Classical and Quantum Computation* (Americal Mathematical Society, 2002).
 - ⁴ S. Das Sarma, M. Friedman and C. Nayak, *Phys. Today*, **59** 32(2006).
 - ⁵ J. Preskill, *Phys. Today*, **52** 24(1999).
 - ⁶ A.Yu. Kitaev, *Annals of Physics*, **321**, (2006), 2-111.
 - ⁷ A.Yu.Kitaev, *Annals of Physics*, **303**, (2003), 2-30.
 - ⁸ G. Baskaran, Saptarshi Mandal, R Shankar, *Phys. Rev. Lett.*, **98**, 247201(2007).
 - ⁹ X. G. Wen, F. Wilczek, and A. Zee, *Phys. Rev. B* **39**, 11413(1989).
 - ¹⁰ N. Read and S. Sachdev, *Phys. Rev. Lett.* **66**, 1773(1991).
 - ¹¹ D. S. Rokhsar and S. A. Kivelson, *Phys. Rev. Lett.* **61**, 2376(1988).
 - ¹² N. Read and B. Chakraborty, *Phys. Rev. B* **40**, 7133(1989).
 - ¹³ R. Moessner and S. L. Sondhi, *Phys. Rev. Lett.* **86**, 1881(2001).
 - ¹⁴ V. Kalmayer and R. B. Laughlin, *Phys. Rev. Lett.* **59**,

- 2095(1987).
- ¹⁵ Senthil. T, Fisher. M.P.A, *Phys. Rev. B* **62**, 7850(2000).
- ¹⁶ Xiao-Yong Feng, Guang-Ming Zhang, Tao Xiang, *Phys. Rev. Lett.* **98**, 087204(2007).
- ¹⁷ Han-Dong Chen and Jiangping Hu, *Phys. Rev. B* **76**, 193101(2007).
- ¹⁸ Han-Dong Chen and Zohar Nussinov, *J. Phys. A: Math. Theor.* **41**, 075001, 2008.
- ¹⁹ Saptarshi Mandal, R Shankar, G Baskaran, Poster 14, International Conference on Mott Insulator, IISc, Bangalore, July,2006,<http://www.physics.iisc.ernet.in/seminars-past.php>.
- ²⁰ K. P. Schmidt, S Dusuel and J. Vidal, *Phys. Rev. Lett.* **100**, 057208(2008).
- ²¹ K. Sengupta, D. Sen, and S. Mondal, *Phys. Rev. Lett.* **100**, 077204(2008).
- ²² G. Baskaran, D. Sen and R. Shankar, *Phys. Rev. B* **78**, 115116(2008).
- ²³ S. Mandal, N Surendran, *Phys. Rev. B* **79**, 024426(2009).
- ²⁴ S. Dusuel, K. P. Schmidt, J. Vidal, *Phys. Rev. Lett.* **100**, 177204(2008).
- ²⁵ G. Baskaran, Z. Zou and P. W. Anderson, *Solid St. Commn.* 63973(1987)
- ²⁶ Ian Affleck, Z. Zou, T. Hsu and P.W. Anderson. *Phys. Rev. B* **38** 745(1988).

- ²⁷ E. Dagotto, E. Fradkin, and A. Moreo. Phys. Rev. B **38** 2926(1988).
- ²⁸ G. Baskaran. *Indian J. Phys.* **80** (6), 583-592(2006).
- ²⁹ G. Baskaran, E. Tosatti and L. Yu, Int. J. Mod. Phys. B, **2**, 555, 5(1988).
- ³⁰ J. B. Martson, Phys. Rev. Lett. **61**, 1914(1988).
- ³¹ E. Fradkin, *Field Theories of Condensed Matter Systems*, HarperCollins Canada / Perseus Books.
- ³² E. Lieb, F. Wu, Phys. Rev. Lett. **20**, 1444(1968).
- ³³ V. Kalmeyer, R. B. Laughlin, Phys. Rev. Lett. **59**, 2095(1987).
- ³⁴ P. W. Anderson, *Science* **235**, 1196(1987).
- ³⁵ P. W. Anderson, *Science* **288**, 480 (2000).
- ³⁶ X. G. Wen, Phys. Rev. B **65**, 165113(2008).
- ³⁷ E. H. Lieb, Phys. Rev. Lett. **73**, 2158 (1994).
- ³⁸ X. G. Wen and Q. Niu, Phys. Rev. B **41**, 9377(1990).
- ³⁹ Abhinav Saket, S. R. Hassan and R. Shankar, Phys. Rev. B **82**, 174409 (2010).
- ⁴⁰ F. J. Burnell and Chetan Nayak, Phys. Rev. B **84**, 125125 (2011).